Chondrex, Inc.

Immunization Grade Chick Type IX Collagen, Lyophilized

Catalog # 1071

For Research Use Only - Not Human or Therapeutic Use

DESCRIPTION:	Type IX collagen purified from pepsin-solubilized chick cartilage by repeat salt precipitation.
	Type IX collagen is one of three types of collagen which make up cartilage fibrils. It consists of three genetically distinct polypeptides, which are cross-linked by disulfide bonds and is digested into two fragments, high molecular weight (HMW) and low molecular weight (LMW) fragments by pepsin digestion from tissues. The ratio of HMW and LMW varies between individual batches.
APPLICATION:	Use as an immunizing antigen to generate antibodies, as an ELISA antigen to detect anti-type IX collagen antibodies and as a standard for gel analysis.
	NOTE: Anti-type IX collagen antibodies which cross-react to autologous type IX collagen may contribute to the induction of polychondritis in certain strains of mice.
QUANTITY:	5 mg
FORM:	Lyophilized powder
SOURCE:	Chick articular cartilage
MOLECULAR WEIGHT:	Intact type IX collagen: approximately 220 kDa by 8% gel analysis under non-reducing conditions.
	Pepsin solubilized type IX collagen is a mixture of three HMW (150, 135, and 120 kDa) and one LMW (35 kDa) fragments.
PURITY:	>90% by SDS-PAGE
STORAGE:	4°C in the dark for lyophilized form and -20°C for solution form. Collagen may gradually degrade under neutral conditions.
STABILITY:	2 years
NOTES:	Type IX collagen can be dissolved at 4 mg/ml in an acidic solution such as 0.01-0.05M acetic acid, pH 3.0-3.3 or 0.15M citrate buffer, pH 3.6 by stirring at 4°C overnight. To neutralize the solution, add 10X neutral buffer containing 1.5M NaCl or dialyze the solution against a neutral buffer.
REFERENCES:	M. Shimokomaki, V. Duance, A. Bailey. FEBS Lett. 121:51-54 (1980)
	C. Reese, H. Wiedemann, K. Kuhn, R. Mayne, Biochemistry. 21:826-830 (1982)
	M. van der Rest, R. Mayne, Y. Ninomiya, et al. J Biol Chem. 260:220-225 (1985)