

## Immunization Grade Bovine Type II Collagen, Solution

Catalog # 20022

*For Research Use Only - Not Human or Therapeutic Use*

DESCRIPTION:	Highly purified type II collagen (treated with 3M guanidine, DEAE-cellulose, and Na <sub>2</sub> HPO <sub>4</sub> )
APPLICATION:	Can be used for various purposes, such as a standard for collagen analysis by SDS-gel, substrate for MMP-1, MMP-8, and MMP-13, and as an antigen for immunizing animals for the induction of collagen-induced arthritis (CIA) in certain species and strains of experimental animals.
QUANTITY:	5 ml
FORM:	2 mg/ml solution in 0.05M acetic acid
SOURCE:	Bovine articular cartilage
MOLECULAR WEIGHT:	300 kDa
PURITY:	>99% by SDS-PAGE, free of pepsin and proteoglycans
STORAGE:	-20°C in the dark
STABILITY:	2 years
NOTES:	<p>The maximum concentration of type II collagen in acidic buffer is 4 mg/ml. In order to neutralize this product, add this collagen solution to a desired neutral buffer such as 2-10X Tris-NaCl (NaCl final concentration: 0.15-0.2M) or dialyze against 1X neutral buffer at 4°C.</p> <p>NOTE: Although type II collagen does not form gel at low concentrations, Chondrex, Inc. recommends keeping the collagen solution on ice.</p>
REFERENCES:	<p>R. Trelstad, <i>et al. Biochemistry</i> <b>9</b>: 4993-4998 (1970)</p> <p>E. Miller, <i>Biochemistry</i> <b>10</b>: 1652-1659 (1971)</p> <p>D. Trentham, <i>et al. J. Exp. Med.</i> <b>146</b>: 857-868 (1977)</p> <p>P. Wooley, <i>et al. J. Immunology</i> <b>135</b>: 2443-2451 (1985)</p>