

Mouse Anti-HDM IgM Antibody Assay Kit

Catalog # 3036

For Research Use Only - Not Human or Therapeutic Use

INTRODUCTION

Asthma is a common chronic inflammatory disease that affects 300 million people of all ages worldwide (1). It is caused by exposure to allergens such as dust mites, pet dander, pollen, or mold, and characterized by airflow obstruction and bronchospasm. House dust mite (HDM) is the most common asthma allergen, which affects up to 85% of asthma patients (2, 3). Of the two main mite species, *Dermatophagoides pteronyssinus* (Der p) and *Dermatophagoides farinae* (Der f), more than 20 types of HDM allergens are defined based on sequential and functional homologies. Among those HDM allergens, group 1 (Der 1) and group 2 (Der 2) dominate overall allergic responses in patients and are the most commonly researched allergens (4-6).

Previously, asthma was considered to be an inflammatory airway disease mediated by the adaptive immune system, particularly type 2 helper T-cells (7). However, recent studies indicate that the innate immune system is also involved in triggering an inflammatory response in both asthma patients and animal models (8-10). These inflammatory changes and airway remodeling significantly vary depending on the types of allergens (11). To meet such needs, a mouse HDM-induced asthma model is a useful tool to dissect the pathological roles of the adaptive and innate immune systems activated by different HDM elements. This is an advantage over the classic ovalbumin-induced asthma model which activates adaptive immunity preferentially.

Recently, it was reported that HDM-specific sublingual immunotherapy (SLIT) is more efficacious than subcutaneous immunotherapy at preventing the development of allergic inflammatory reactions in a mouse model (12). This SLIT protocol has been approved as a treatment to reduce allergy or asthma symptoms in patients (13).

To study the immune response to allergens and allergen-specific pathological effects in mouse models, Chondrex, Inc. provides mouse ELISA kits for assaying anti-HDM IgG, IgM, and IgE antibodies (Catalog # 3030, 3036, 3037), in addition to anti-HDM antibody IgG subclasses IgG1, IgG2a, IgG2b, and IgG3 (Catalog # 3034, 3038, 3035, 3039). Chondrex, Inc. also offers ELISA kits for assaying anti-OVA antibody subtypes IgA, IgE, IgG, and IgM and anti-OVA IgG subclasses IgG1, IgG2a, IgG2b, and IgG2c, (Catalog # 3004, 3010, 3011, 3015 - 3018, 3029) as well as total immunoglobulin subtypes IgA, IgE, IgG, and IgM, and IgG subclasses IgG1, IgG2a, IgG2b, and IgG3, (Catalog # 3005, 3019, 3023 - 3028).

Note: Other antibody subtype ELISA kits against HDM as well as HDM antigen detection kits are currently under development. Please contact Chondrex, Inc. support (support@chondrex.com) for more information.

KIT COMPONENTS


Item	Quantity	Amount	Storage
Standard Anti-HDM Mouse IgM Antibody (30361)	1 vial	1000 ng/vial, lyophilized	-20°C
Secondary Antibody - Peroxidase-Conjugated Goat Anti-Mouse IgM Antibody (30363)	2 vials	50 µl	-20°C
Solution B - Blocking Buffer (30313)	1 bottle	10 ml	-20°C
Solution C - Sample/Standard/Detection Antibody Dilution Buffer (30314)	1 bottle	50 ml	-20°C
TMB Solution (90023)	2 vials	0.2 ml	-20°C
Chromogen Dilution Buffer (90022)	1 bottle	20 ml	-20°C
Stop Solution - 2N Sulfuric Acid (9016)	1 bottle	10 ml	-20°C
Wash Buffer, 20X (9005)	1 bottle	50 ml	-20°C
HDM from <i>Dermatophagoides pteronyssinus</i> Coated ELISA Plate	1 each	96-well (Brown 8-well strips x 12)	-20°C

NOTES BEFORE USING ASSAY

- Note 1: It is recommended that the standard and samples be run in duplicate.
- Note 2: Warm up all buffers to room temperature before use.
- Note 3: Partially used reagents may be kept at -20°C .
- Note 4: Crystals may form in the 20X wash buffer when stored at cold temperatures. If crystals have formed, warm the wash buffer by placing the bottle in warm water until crystals are dissolved completely.
- Note 5: Measure exact volume of buffers using a serological pipette, as extra buffer is provided.
- Note 6: Cover the plate with plastic wrap or a plate sealer after each step to avoid the edge effect.

ASSAY PROCEDURE


- Dilute Wash Buffer:** Dilute 50 ml of 20X wash buffer in 950 ml of distilled water (1X wash buffer).
- Add Blocking Buffer:** Add 100 μl of the Blocking Buffer (Solution B) to each well and incubate for 1 hour at room temperature.
- Prepare Standard Dilutions:** The recommended standard range is 16-1000 ng/ml. Dissolve one vial of Standard (1000 ng/vial) in 1 ml of Sample/Standard/Detection Antibody Dilution Buffer (Solution C) and keep it as a 1000 ng/ml standard stock. Add 250 μl of this standard solution to 250 μl of Solution C to make a 500 ng/ml solution. Then, serially dilute it with Solution C. For example, mix 250 μl of the 500 ng/ml solution with an equal volume of Solution C to make a 250 ng/ml solution, and then repeat it five more times for 125, 63, 31, and 16 ng/ml standard solutions.



- Prepare Sample Dilutions:** The dilution of serum from mouse immunized with HDM varies (1:10 or more) depending on the immunization schedule and timing of serum collection. In general, no IgM antibodies against HDM are observed in normal serum at a 1:10 dilution.
- Wash:** Wash the plate with 1X wash buffer at least 3 times using a wash bottle with manifold or an automated plate washer. Empty the plate by inverting it and blot on a paper towel to remove excess liquid. *Do not allow the plate to dry out.*

6. **Add Standards and Samples:** Add 100 μ l of standards, Solution C (blank), and samples to wells in duplicate. Incubate at room temperature for 2 hours.

Figure 1 - Mouse anti-HDM IgM Assay Standard layout.



7. **Wash:** Wash the plate with 1X wash buffer at least 3 times using a wash bottle with manifold or an automated plate washer. Empty the plate by inverting it and blot on a paper towel to remove excess liquid. *Do not allow the plate to dry out.*
8. **Add Secondary Antibody:** Dissolve one vial of Secondary Antibody in 10 ml Sample/Standard/Detection Antibody Dilution Buffer (Solution C). Add 100 μ l of detection antibody solution to each well and incubate at room temperature for 1 hour.
9. **Wash:** Wash the plate with 1X wash buffer at least 3 times using a wash bottle with manifold or an automated plate washer. Empty the plate by inverting it and blot on a paper towel to remove excess liquid. *Do not allow the plate to dry out.*
10. **Add TMB:** Use new tubes when preparing TMB. Dilute one vial of TMB with 10 ml Chromogen Dilution Buffer just prior to use. Add 100 μ l of TMB solution to all wells immediately after washing the plate and incubate for 25 minutes at room temperature.
11. **Stop:** Add 50 μ l of 2N sulfuric acid (Stop Solution) to each well.
12. **Read Plate:** Read the OD values at 450 nm (a 630 nm filter can be used as a reference). If the OD values of samples are greater than the OD values of the highest standard, re-assay the samples at a higher dilution.

CALCULATION OF ANTIBODY TITERS

1. Average the duplicate OD values for the standards, blanks (B), and test samples.
2. Subtract the averaged blank (B) values from the averaged OD values of the standards and test samples.
3. Plot the OD values of the standards against the ng/ml of antibody standard. Using a log/log plot will linearize the data. Figure 2 shows an example of a standard curve of anti-HDM IgM antibodies.
4. The ng/ml of antibody in test samples can be calculated using regression analysis. Multiply it by the sample dilution factor to obtain the antibody concentration (ng/ml) in original sample specimens.

Figure 2 - A typical standard curve for anti-HDM IgM assay




Table 1 - Reproducibility of data assayed by Mouse Anti-HDM IgM Antibody Assay Kit

Test At	32 ng/ml	125 ng/ml	500 ng/ml
Inter-Assay CV (%)	3.8	4.8	0.9
Intra-Assay CV (%)	8.3	4.0	3.1
Spiking Test*	103%	105%	102%

Known amounts of anti-HDM IgM were added to samples and then diluted with Sample/Standard/Detection Antibody Dilution Buffer to assay anti-HDM IgM antibodies by ELISA.

REFERENCES

1. M. Masoli, D. Fabian, S. Holt, R. Beasley, G. I. f. A. G. Program, The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy* 59, 469-478 (2004).
2. L. G. Gregory, C. M. Lloyd, Orchestrating house dust mite-associated allergy in the lung. *Trends Immunol* 32, 402-411 (2011).
3. V. D. Gandhi, C. Davidson, M. Asaduzzaman, D. Nahirney, H. Vliagoftis, House dust mite interactions with airway epithelium: role in allergic airway inflammation. *Curr Allergy Asthma Rep* 13, 262-270 (2013).
4. W. R. Thomas, W. Smith, House-dust-mite allergens. *Allergy* 53, 821-832 (1998).
5. A. Custovic, S. C. Taggart, H. C. Francis, M. D. Chapman, A. Woodcock, Exposure to house dust mite allergens and the clinical activity of asthma. *J Allergy Clin Immunol* 98, 64-72 (1996).
6. A. Jacquet, The role of innate immunity activation in house dust mite allergy. *Trends Mol Med* 17, 604-611 (2011).
7. L. Cohn, J. A. Elias, G. L. Chupp, Asthma: mechanisms of disease persistence and progression. *Annu Rev Immunol* 22, 789-815 (2004).
8. J. R. Murdoch, C. M. Lloyd, Chronic inflammation and asthma. *Mutat Res* 690, 24-39 (2010).
9. J. C. Virchow et al., Efficacy of a House Dust Mite Sublingual Allergen Immunotherapy Tablet in Adults With Allergic Asthma: A Randomized Clinical Trial. *JAMA* 315, 1715-1725 (2016).
10. S. Hagner et al., House Dust Mite-Specific Sublingual Immunotherapy Prevents the Development of Allergic Inflammation in a Mouse Model of Experimental Asthma. *Int Arch Allergy Immunol* 170, 22-34 (2016).
11. C. S. Stevenson, M. A. Birrell, Moving towards a new generation of animal models for asthma and COPD with improved clinical relevance. *Pharmacol Ther* 130, 93-105 (2011).
12. P. Moingeon et al., Immune mechanisms of allergen-specific sublingual immunotherapy. *Allergy* 61, 151-165 (2006).
13. Z. Aryan, E. Compalati, E. Comapalati, G. W. Canonica, N. Rezaei, Allergen-specific immunotherapy in asthmatic children: from the basis to clinical applications. *Expert Rev Vaccines* 12, 639-659 (2013).