

# FITC-Polysucrose


**Trade name:** FITC-Polysucrose

**Chemical names:** Polysucrose (3',6'dihydroxy3-oxospiro (isobenzofuran(3H),9'[9H] xanthen]-5(or 6)-yl) carbamothioate

Fluorescein isothiocyanate/Fluoresceinyl thiocarbamoyl

Polysucrose/FITC-Ficoll®

**CAS nr:** N/A



*Fig. 1 Structural representation of fragment of FITC-Polysucrose molecule.*

## Synthesis

Polysucrose fractions are labelled with fluorescein by a procedure similar to that described by de Belder and Granath (1). The fluorescein moiety is attached by a stable thiocarbamoyl linkage and the labelling procedure does not lead to any depolymerisation of the Polysucrose. The FITC-Polysucroses carry from 0.001-0.008 mol. FITC per glucose unit and at these low levels of substitution confer minimal charges to the Polysucrose, which is an essential requirement for permeability studies.

## Description

FITC-Polysucrose is supplied as a yellow powder which dissolves freely in water or salt solutions giving a yellow solution. The product also dissolves in DMSO, formamide and certain other polar organic solvents but is insoluble in lower aliphatic alcohols, acetone, chloroform, dimethylformamide.

Polysucrose (Ficoll®) is synthesized by polymerizing sucrose with epichlorohydrin. The resulting polymer is highly branched and behaves as a globular molecule in solution (see section Physical chemical properties). Polysucrose contains only primary and secondary hydroxyl groups.

## Spectral data

Excitation is best performed at 496nm and fluorescence measured at 525 nm (see Fig.2). The dependence of fluorescence from a FITC-Polysucrose solution in the range pH 3-9 is shown in Fig.2. Measurements in biological media may significantly affect the fluorescence intensity which may be enhanced or depressed.

## Physical chemical properties

The Polysucrose molecule behaves as a globular molecule in solution as is to be expected from its structure. In Table 1 (below) a comparison of the Stokes radius of dextran and Polysucrose fractions reflects these differences in molecular flexibility. The molecule is best regarded as an intermediate between a hard-solid sphere and a flexible coil. Thus, when comparing Polysucrose and dextran fractions of similar molecular weights, the molecular dimensions of the Polysucrose will always be smaller. It is unsuitable to use dextran calibration for GPC determinations of the molecular weight of Polysucrose products. Polysucrose solutions have very low osmotic pressures compared to sucrose solution of equivalent concentrations. Thus a 10% solution of Polysucrose 70 has an osmolality of 3 mOs/kg compared to 150 for a 10% sucrose.

MW *10 <sup>3</sup>	Dextran Stokes radius	Polysucrose Stokes radius	Albumin Stokes radius
500	147	106	-
70	58	49.5	35
49	44.5	40	-

*Table 1. Molecular dimensions of Polysucrose and dextran expressed as Stokes Radii (Å)*

## Storage and stability

FITC-Polysucrose powder when stored in airtight containers at ambient temperatures is stable for at least 6 years in solution. The stability of FITC-Polysucroses has not been investigated in detail. However, the stability of the thiocarbamoyl linkage between the fluorescein moiety and Polysucrose will be similar to that with dextran (see data-file for information on stability of FITC-dextran). Only at elevated pH (>9) and elevated temperatures is there a risk for hydrolysis of the thiocarbamoyl linkage.

FITC-dextran was found to be stable at pH4 for up to 1 month at temperatures up to 35°C but this is not to be recommended for Polysucrose based products owing to the lability of the glycosidic linkages in sucrose. Polysucrose itself can be autoclaved at neutral and slightly alkaline pH.

## Toxicity

Polysucrose fractions exhibit no toxic symptoms when tested orally or intravenously. Intravenously Polysucrose fractions (100 000 to 500 000) when administered at doses up to 12 g/kg in experimental animals showed no toxic symptoms. Polysucrose is not however degraded in the blood and accumulates in the liver, spleen and kidneys. Polysucrose shows excellent biocompatibility with cells, virus or microorganisms and has been used for many decades in separation technology.

## Biological Aspects and Applications

Polysucrose (Ficoll®) and derivatives thereof offer many interesting characteristics for studying the physiology of various organs. Many articles on the glomerular membrane have appeared over the years. Polysucrose unlike dextran has a more compact globular structure but nevertheless appears to possess some flexibility and is best regarded as an intermediate between a hard-globular protein and the loose flexible coil of dextran. Polysucroses exhibit excellent biocompatibility and are not secreted or reabsorbed by the renal tubules.

The diffusion coefficients of narrow fractions of inter alia dextran and Polysucrose has been studied in membranes (2). It was shown that the diffusion properties of Polysucrose differed significantly from the linear polymers which diffused faster through the membranes. The authors concluded that Polysucrose behaves more like a solid sphere. A close examination of the size and conformation of Polysucrose using GPC in combination with light-scattering and viscosity detectors indicated that the Polysucrose molecule is best regarded as intermediate between a solid sphere and a well-solvated linear random coil (3). Venturoli and Rippe (4) have reviewed the available data on the two polysaccharides Ficoll and dextran for assessing the glomerular perm selectivity as compared to glomerular proteins. The authors elaborate on the various properties which may influence the results such as molecular size, shape, charge and flexibility and assess their results in various pore models.

FITC-Polysucroses (FITC-Ficoll®) are primarily used for studying permeability and transport in cells and vessels and tissues. An added benefit is that measurements of the fluorescence provide quantitative data on transport and permeability of healthy and diseased tissues. Such studies can be performed in real time by intravital fluorescence microscopy. The technique offers high sensitivity and concentrations down to 1µg/ml can be detected in tissue fluids.

## General procedures

The microvasculature of the hamster cheek pouch has proved to be a useful model for studying plasma leakage in different experimental conditions, e.g. following ischemia/reperfusion, topical application of a whole range of inflammatory mediators, parasites and bacteria. With this technique, vascular permeability changes can be studied in real time and be related to other microvascular events such as leukocyte adhesion and activation. The cheek pouches are examined by intravital fluorescence microscopy using suitable filters (490/520nm) and images are captured with a digital camera. A suitable concentration of FITC-Polysucrose for infusion in experimental animals is 5 % (approx. 100mg/kg bodyweight) (5-7). An alternative procedure using rabbit ear chambers has been described. The regenerative titanium ear-chambers (rabbits) were used to study the blood/lymph systems in the microcirculation with fluorescent-dextran. Lymph ingrowth is seen after 4-8 weeks of implantation (8).

## 1. Permeability studies in cells

The effect of LPS on different splenic non-lymphoid cells was monitored by selective uptake of FITC-Polysucrose (9). Marginal zone macrophages could be distinguished by the lack of uptake of FITC-Polysucrose and antibody staining (10). Other studies on liposomes and non-lymphoid cells using FITC-Polysucroses have been reported (11,12).

## 2. Permeability studies on the glomerular membrane

The available data on the two polysaccharides Polysucrose (Ficoll®) and dextran for assessing the glomerular perm selectivity as compared to glomerular proteins has been reviewed (13). Polydisperse polysaccharides are excellent probes for measuring glomerular perm selectivity and are reproducible, reliable and elegant. The authors elaborate on the various properties which may influence the results such as molecular size, shape, charge and flexibility and assess their results in various pore models.

Studies of the glomerular permeability of Polysucrose when infused intravenously showed that it has a cut-off at about 50Å whereas dextran is excreted up to 60-70Å – this is explained by the greater flexibility of dextran. The clearance of FITC-Polysucrose in mice lacking endothelial caveolae was studied in order to elucidate macromolecular transport pathways (14). The glomerular filter was studied at different glomerular filtration rates using FITC-Polysucrose 70 and 400 (also FITC-inulin) (15).

Studies on the glomerular filtration of dextran and Polysucrose showed that the glomerular membrane presented a much more restrictive barrier to Polysucrose than to dextran (16).

Interestingly, the values of the sieving coefficient  $\theta$  for Polysucrose approximated to those reported for uncharged globular proteins. Glomerular sieving in rats following surgery and muscle trauma monitored using FITC-Polysucrose 70/400(17). The rats were dosed with a mixture of FITC-Polysucrose 400 (960µg), FITC-Polysucrose 70 (40µg) and FITC-inulin (500µg) as a priming bolus. Glomerular sieving measured in caveolin-1 knockout mice monitored using FITC-Polysucrose 70/400 was used to investigate the factors affecting glomerular permeability (18).

FITC-Polysucrose 70 and albumin were used to estimate the fractional clearances in mice before and after treatment with enzymes degrading various glycosaminoglycans in the glycocalyx (19). FITC-Polysucrose 70 and albumin were infused in rats to explore the effects of temperature and ammonium chloride on the fractional clearance. The sieving coefficients between 8 and 37°C were not significantly different. Polysucroses perform differently to dextrans and theta was lower over the range 20 -70 Å than the corresponding dextrans. Solute shape and may outweigh size and charge (20,21).

To further explain the transport of protein across the capillary walls, mice lacking endothelial caveolae were studied with various permeability probes including FITC-Polysucrose (22).

Fractional clearance of FITC-Polysucrose was determined at low ionic strength in perfusion fixed isolated kidney (23). Perfusates contained approx. 70mg FITC-Polysucrose 70/L were used to evaluate whether the increase in clearance of native albumin after 9 weeks of diabetes was due to reduced charge selectivity or to an alteration in the proportion of large pores (24).

More recent studies have explored the role of nitric oxide (25), reactive oxygen species (26) and scavengers(27) on glomerular permeability.

## List of references

1. A.N.de Belder and K.Granath. Preparation and properties of fluorescein labelled dextrans. *Carbohydr Res.* 1973; 30:375-378
2. G.D.Davidson and W.M.Deen. Hindered diffusion of water-soluble molecules in membranes. *Macromolecules.*1988;21:3474-3481
3. W.H.Fissell, C.L.Hofmann, and R.Smith. Size and conformation of Ficoll as determined by size-exclusion chromatography followed by multiangle light scattering. *Am J Physiol Renal Physiol.* 2010;298: F205-8
4. D.Venturoli and B.Rippe. Polysucrose and dextran vs. globular proteins as probes for testing glomerular permselectivity effects of molecular size, shape, charge and deformability. *Am J Physiol Renal Physiol.* 2005; 25:77-4
5. D.Hultström and E.Svensjö. Intravital and electron microscopy study of bradykinin induced vascular permeability changes using FITC-Ficoll as a tracer. *Journal of Pathology.*1979;129:125-133
6. E.Svensjö. The hamster cheek pouch as a research model in inflammation. Chapter 30 In: David Shepro (Editor), *Microvascular Research – Biology and Pathology.* p.195-207, 2006. Elsevier Academic Press.
7. E.Svensjö, E.M.Saraiva, M.T.Bozza et al. Salivary gland homogenates of *Lutzomyia longipalpis* and its vasodilatory peptide maxadilan cause plasma leakage via PAC1 receptor activation. *J.Vasc.Res.* 2009;46:435446
8. J.Jonsson, K.E.Arfor, and H.C.Hint. Studies on relationships between blood and lymphatic systems within the microcirculation. 6th Europ.Conf.Microcirculation, Aalborg. 1970;214-218 (Karger, Basel 1971)
9. P.L.Amlot, D.Grennan and J.H.Humphrey. Splenic dependence of the antibody response to thyroid independent (TI-2) antigens. *Eur J Immunol.* 1985; 15:50812
10. G.Kraal and M. Janse. Marginal metallophilic cells of the mouse spleen identified by a monoclonal antibody. *Immunology.*18;58:665-9
11. P.H.Groeneveld, T.Erich and G.Kraal. The differential effects of Bacterial lipopolysaccharides (LPS) on splenic non-lympoid cells demonstrated by monoclonal antibodies. *Immunology.* 1986; 58:285-90
12. E.Claassen. Post-formation fluorescent labelling of liposomal membranes. In vivo detection, localization and kinetics. *J Immuno Methods.* 1992; 147:231-40
13. D.Venturoli and B.Rippe. Ficoll and dextran vs. Globular proteins as probes for glomerular permselectivity; effects of molecular size, shape, charge and deformability. *Am J Physiol Renal Physiol.* 2005; 25:7784
14. B.I.Rosengren, A.Rippe, C.Rippe et al. Transvascular protein transport in mice endothelial caveolae. *Am J Physiol Heart Circ Physiol.* 2006;291:H1371-7
15. C. Rippe, D. Asgeirsson, D.Venturoli et al. Effects of glomerular filtration rate on the Ficoll sieving coefficients in rats. *Kidney.* 2006; 69:1326-32
16. J.D.Oliver III, S.Andersson, J.L.Troy et al. Determination of glomerular size-selectivity in the normal rat with Ficoll. *J Amer Soc Nephrol.* 1992; 3:214-22
17. J.Axelsson, I.Mahmutovic, A.Rippe et al. Loss of size selectivity of the glomerular filtration barrier in rats following laparotomy and muscle trauma. *J Physiol Renal Physiol.* 2009;297: F577-82
18. G.Grände, C.Rippe, A.Rippe et al. Unaltered size selectivity of the glomerular filtration in caveolin-1 knockout mice. *Am J Physiol Renal Physiol.* 2009;297: F257-62

19. M.Jeansson and B.Haraldsson. Glomerular size and charge selectivity in the mouse after exposure to glucosaminoglycan degrading enzymes *J Am Soc Nephrol.* 2003; 14:1756-65
20. M.Ohlsson, J.Sörensson and B.Haraldsson, Glomerular size and charge selectivity in the rat as revealed by FITC-Ficoll and albumin. *Am J Physiol Renal Physiol.*
21. M.Ohlsson, J.Sörensson and B.Haraldsson. Glomerular size and charge selectivity in the rat as revealed by FITC-Ficoll and albumin. *Am J Physiol Renal Physiol.* 2001;278:F992-3.1986; 58: 655-9.
22. B-I.Rosengren, A.Rippe, C.Rippe et al. Transvascular protein transport in mice lacking endothelial caveolae. *Am J Physiol Heart Circ Physiol.* 2006;291:H1371-7
23. G.Ciarimboli, C.Hjalmarsson, LBokenkamp et al. Dynamic alterations of glomerular charge density in fixed rat kidneys suggest involvement of endothelial coat. *Am J Physiol Renal Physiol.* 2003;285: F722-30
24. C.Rippe, A.Rippe, O.Torffvit et al. Size and charge selectivity of the glomerular filter in early experimental diabetes in rats. *Am Physiol Renal Physiol.* 2007;293:F1533-8
25. J.Dolinina, K.Sverrisson, A.Rippe, et al. Nitric oxide synthetase inhibition causes acute increases in glomerular permeability in vivo, dependent upon reactive oxygen species (ROS). *Am J Physiol Renal Physiol.* 2016 Sept 28;ajprenal.00152.2016.
26. K.Sverrisson J.Axelsson, A.Rippe et al. Acute reactive oxygen species (ROS)-dependent effects of IL-1 $\beta$ , TNF- $\alpha$ , and IL-6 on the glomerular filtration barrier (GFB) in vivo. *Am J Physiol Renal Physiol.* 2015 Nov 1;309(9):F800-6
27. J.Axelsson, A.Rippe, K. Sverrisson and B.Rippe. Scavengers of reactive oxygen species, paracalcitol, RhoA, and Rac-1 inhibitors and tacrolimus inhibit angiotensin II-induced actions on glomerular permeability. *Am J Physiol Renal Physiol.* 2013 Aug 1;305(3):F237-43