



ECM Analysis: Solubilization Guide



Analysis of Collagen, Glycosaminoglycans, and DNA

Cartilage tissue engineering generally requires the analysis of collagen, glycosaminoglycans (GAGs), and DNA in conjunction with cellular or extracellular matrix components (ECM) (1, 2). Consequently, depending on the analytes, appropriate sample preparation protocols must be utilized. Table 1 shows three sample preparation protocols and assay kits as guidelines for ECM analysis. Please contact Chondrex, Inc. at support@chondrex.com for more information.

- A. Guanidine hydrochloride (GuCl) extracts GAGs, but fails to completely extract collagen and DNA, resulting in underestimated values (1, 3).
- B. Papain digestion solubilizes GAGs (4) and DNA (5) but degrades collagen.
- C. Chondrex, Inc.'s collagen solubilization protocol can be used on samples for the GAGs assay (6), but not for the DNA assay.

Protocol	Analytes				
	Target	Sulfated GAGs	DNA	Total Collagen	Individual Native Collagen Types
A) Guanidine Hydrochloride (GuCl) Extraction*	Yes	Yes*	No		No
B) Papain Digestion	Yes	Yes	Yes		No
C) GuCl Extraction*, Pepsin Digestion, and Elastase Digestion	Yes	No	Yes		Yes
Recommended Assays (Catalog #)	6022	6023	6017	6012-16, 6018, 6019, 6021	

*GuCl extraction works for samples consisting of chitosan or agarose, but not for tissues such as cartilage due to poor extraction efficiency (60%) (1).

References

1. C. D. Hoemann, J. Sun, V. Chrzanowski, M. D. Buschmann. A multivalent assay to detect glycosaminoglycan, protein, collagen, RNA, and DNA content in milligram samples of cartilage or hydrogel-based repair cartilage. *Anal Biochem* **300**, 1-10 (2002).
2. D. J. Griffon, M. R. Sedighi, D. V. Schaeffer, J. A. Eurell, A. L. Johnson. Chitosan scaffolds: interconnective pore size and cartilage engineering. *Acta Biomater* **2**, 313-320 (2006).
3. S. W. Sajdera, V. C. Hascall. Protein polysaccharide complex from bovine nasal cartilage. A comparison of low and high shear extraction procedures. *J Biol Chem* **244**, 77-87 (1969).
4. R. W. Farndale, D. J. Buttle, A. J. Barrett. Improved quantitation and discrimination of sulphated glycosaminoglycans by use of dimethylmethylene blue. *Biochem Biophys Acta* **883**, 173-177 (1986).
5. Y. J. Kim, R. L. Sah, J. Y. Doong, A. J. Grodzinsky. Fluorometric assay of DNA in cartilage explants using Hoechst 33258. *Anal Biochem* **174**, 168-176 (1988).
6. K. von der Mark, M. van Menzel, H. Wiedemann. Isolation and characterization of new collagens from chick cartilage. *Eur J Biochem* **124**, 57-62 (1982).