Complete Freund's Adjuvant

Catalog # 7008

For Research Use Only - Not Human or Therapeutic Use

DESCRIPTION:	Heat-killed *M. Tuberculosis* H37 RA (non-viable) suspended at a concentration of 1 mg/ml in incomplete Freund's adjuvant
APPLICATION:	Use as an adjuvant to induce collagen-induced arthritis (CIA) in mice and to induce adjuvant-induced arthritis (AIA) in rats
QUANTITY:	5 ml
FORM:	1 mg/ml solution
EMULSION:	Emulsify with an equal volume of collagen using an electric homogenizer. The syringe-syringe method is not recommended to make collagen-CFA emulsions for inducing arthritis, but it can be used for obtaining anti-sera.
STORAGE:	4ºC in the dark
STABILITY:	1 year
NOTES:	Before use, please vortex well to ensure consistent *M. tuberculosis* dosage
REFERENCES:	C. William, *et al. J. Exp. Med*. **162**: 1878-1891 (1985)
	I. Campbell, *et al. Eur. J. Immunol.* **30**: 1568-1575 (2000)

© 2020 Chondrex, Inc. All Rights Reserved, 7008 4.0

16928 Woodinville-Redmond Rd NE Suite B-101
Woodinville, WA 98072

Phone: 425.702.6365 or 888.246.6373
Fax: 425.882.3094

www.chondrex.com
support@chondrex.com