Immunization Grade Bovine Type II Collagen, Solution

Catalog # 20022

For Research Use Only - Not Human or Therapeutic Use

DESCRIPTION:	Highly purified type II collagen (treated with 3M guanidine, DEAE-cellulose, and Na2HPO4)
APPLICATION:	Can be used for various purposes, such as a standard for collagen analysis by SDS-gel, substrate for MMP-1, MMP-8, and MMP-13, and as an antigen for immunizing animals for the induction of collagen-induced arthritis (CIA) in certain species and strains of experimental animals.
QUANTITY:	5 ml
FORM:	2 mg/ml solution in 0.05M acetic acid
SOURCE:	Bovine articular cartilage
MOLECULAR WEIGHT:	300 kDa
PURITY:	>99% by SDS-PAGE, free of pepsin and proteoglycans
STORAGE:	-20°C in the dark
STABILITY:	2 years
NOTES:	The maximum concentration of type II collagen in acidic buffer is 4 mg/ml. In order to neutralize this product, add this collagen solution to a desired neutral buffer such as 2-10X Tris-NaCl (NaCl final concentration: 0.15-0.2M) or dialyze against 1X neutral buffer at 4°C. NOTE: Although type II collagen does not form gel at low concentrations, Chondrex, Inc. recommends keeping the collagen solution on ice.
REFERENCES:	R. Trelstad, *et al. Biochemistry* **9**: 4993-4998 (1970) E. Miller, *Biochemistry* **10**: 1652-1659 (1971) D. Trentham, *et al. J. Exp. Med*. **146**: 857-868 (1977) P. Wooley, *et al. J. Immunology* **135**: 2443-2451 (1985)

© 2020 Chondrex, Inc. All Rights Reserved, 20022 2.0

16928 Woodinville-Redmond Rd NE Suite B-101
Woodinville, WA 98072

Phone: 425.702.6365 or 888.246.6373
Fax: 425.882.3094

www.chondrex.com
support@chondrex.com